



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

404. Proposed by S. LEFSCHETZ, Ph. D., University of Nebraska.

Let ABC be a triangle, M a point on BC , BE the intersection of the perpendicular to AM in M with AB and AC , F the other intersection of the circle circumscribed to ABC with the circle through M and A orthogonal to it in A . Prove that the points C, D, E, F are on a circle, and find the envelope of the latter when M describes BC .

No solution of this problem has been received.

405. Proposed by C. N. SCHMALL, New York City.

A plane cuts a constant volume from a given right cone. Prove that the minor axis of the section has a constant length.

Solution by the PROPOSER.

Let C be the vertex of the cone, AB the major axis of any one of the sections made by the plane. Let F be the point of contact of the inscribed circle of triangle ABC with the side AB . Then F is a focus of the section. Draw CD perpendicular to AB . Let V denote the volume cut off. Then, if α, β , denote the semi-axes of the section, we have

$$\begin{aligned} V &= \frac{1}{3} CD \cdot (\text{area of section}) = \frac{1}{3} CD \cdot \pi \alpha \beta = \frac{1}{3} \pi \cdot CD \cdot \alpha \beta \\ &= \frac{1}{6} \pi \cdot CD \cdot AB \cdot \beta = \frac{1}{6} \pi ab \sin C \cdot \beta \dots (1), \end{aligned}$$

(where a, b , are the sides AC, BC , of the triangle ABC). But, $\beta^2 = AF \cdot FB = (s-a)(s-b) = ab \sin^2 \frac{1}{2} C$. $\therefore ab = \beta^2 \cos^2 \frac{1}{2} C$.

Substituting this in (1), we get, $V = \frac{1}{6} \pi \beta^3 \cdot \tan \frac{1}{2} C$.

Now, since C and V are constant, therefore β , or the minor axis (2β) is also constant. It is now also evident, by (1), that the area of triangle ABC is constant.

406. Proposed by DR. R. K. MORLEY, University of Illinois.

Given the lengths of a pair of conjugate diameters of an ellipse and the angle between them; to construct (with ruler and compass) the axes of the ellipse, *i. e.*, find their lengths and the angles they make with the given diameters.

I. Solution by A. M. HARDING, University of Arkansas.

Let PP' and QQ' be the conjugate diameters, and C the center. Produce CQ to K , so that $CQ \cdot QK = CP^2$.

Draw a line through Q parallel to CP . Construct a circle with its center on this line and passing through C and K . Let the circle cut the line through Q at M and N . Then the axes of the ellipse will lie along CM and CN .